

## Astronomy 310 — Final Exam

1. The parallax of a star is 10 milli-arcseconds, and its apparent visual magnitude is 3.5. What is its absolute visual magnitude?
  - (a) 5.5
  - (b) 3.5
  - (c) 1.5
  - (d) -1.5 [\*]
  - (e) -3.5
2. A star has a radius of  $3 R_{\odot}$ , and an effective temperature twice that of the Sun. What is its luminosity?
  - (a)  $1 L_{\odot}$
  - (b)  $6 L_{\odot}$
  - (c)  $49 L_{\odot}$
  - (d)  $144 L_{\odot}$  [\*]
  - (e)  $256 L_{\odot}$
3. Consider a binary system where the semi-major axis shrinks by a factor of two, due to tidal interactions. What is the change in the period (expressed as the ratio of the new period to the old period)?
  - (a)  $\frac{1}{2^{3/2}}$  [\*]
  - (b)  $\frac{1}{2^{2/3}}$
  - (c)  $\frac{1}{2}$
  - (d)  $\frac{1}{2^3}$
  - (e)  $\frac{1}{3^2}$
4. In which sort of binary system can the stellar radii be determined from the light curve?
  - (a) Eclipsing [\*]
  - (b) Spectroscopic
  - (c) Visual
  - (d) Spectrum
  - (e) Optical
5. What is the correct (temperature) ordering of the Harvard spectral classification system?
  - (a) A-B-F-G-K-M-O
  - (b) F-B-A-M-O-G-K
  - (c) O-G-K-A-B-M-F
  - (d) O-B-A-F-G-K-M [\*]
  - (e) K-F-O-G-M-A-B
6. What equation can be used to calculate the relative number of atoms in each ionization stage of a given element?
  - (a) Boltzmann
  - (b) Saha [\*]
  - (c) Fermi-Dirac
  - (d) Planck
  - (e) Bose-Einstein

7. In terms of the specific intensity  $I$ , what is the correct expression for the amount of radiation flowing per unit time, per unit solid angle through a unit area at an angle  $\theta$  to the normal?
  - (a)  $I \sin \theta$
  - (b)  $I \theta$
  - (c)  $I \cos \theta$  [\*]
  - (d)  $I \cos \theta \sin \theta$
  - (e)  $I$
8. A beam of radiation with specific intensity  $I$  is normally incident on an absorbing slab with optical thickness  $\tau$ . What is the specific intensity of the radiation coming out of the slab?
  - (a)  $I$
  - (b)  $I \tau$
  - (c)  $I \log(\tau)$
  - (d)  $I \exp(\tau)$
  - (e)  $I \exp(-\tau)$  [\*]
9. Which of these opacity sources does not depend on wavelength?
  - (a) Bound-free
  - (b) Bound-bound
  - (c) Electron scattering [\*]
  - (d) Free-free
  - (e)  $H^-$
10. Which of these mechanisms is responsible for the broad line profiles seen in white dwarf stars?
  - (a) Thermal Doppler broadening
  - (b) Pressure/Collisional broadening [\*]
  - (c) Natural broadening
  - (d) Turbulent broadening
  - (e) Rotation
11. What is the curve of growth?
  - (a) The increase in the radius  $R$  of a star as its mass  $M$  is increased.
  - (b) The increase in the number of ionized atoms in an atmosphere as the effective temperature  $T_{\text{eff}}$  is increased.
  - (c) The increase in the mass  $M$  of a star as it accretes from a binary companion.
  - (d) The increase in the core radius  $R$  of a star as it burns fuel.
  - (e) The increase of the equivalent width  $W$  of a line profile as the column density  $N$  of absorbers is increased. [\*]
12. Over what timescale does a star respond to departures from hydrostatic equilibrium?
  - (a) Virial
  - (b) Nuclear
  - (c) Kelvin-Helmholtz
  - (d) Dynamical [\*]
  - (e) Thermal
13. According to the virial theorem, if a star without any nuclear reactions contracts, what must happen to the thermal energy?

- (a) It must stay the same
  - (b) It must decrease by the change in gravitational energy
  - (c) It must increase by the change in gravitational energy
  - (d) It must decrease by half the change in gravitational energy
  - (e) It must increase by half the change in gravitational energy [\*]
14. If the pressure everywhere in the star follows the relation  $P \propto \rho^\gamma$ , for some arbitrary constant  $\gamma$ , what sort of structure does the star have?
- (a) Isothermal
  - (b) Adiabatic
  - (c) Isobaric
  - (d) Polytropic [\*]
  - (e) Degenerate
15. Which of these mass-fraction combinations corresponds most closely to Solar abundance?
- (a)  $X = 0.7, Z = 0.02$  [\*]
  - (b)  $X = 0.7, Z = 0.0001$
  - (c)  $X = 0.9, Z = 0.01$
  - (d)  $X = 0.3, Z = 0.002$
  - (e)  $X = 0.4, Z = 0.4$
16. In a star in thermal equilibrium, what must be true if  $L_r$  is locally constant (i.e., doesn't vary with radius)?
- (a)  $M_r$  is constant
  - (b)  $\epsilon$  is zero [\*]
  - (c)  $T$  decreases outward
  - (d)  $P$  decreases outward
  - (e)  $\nabla = \nabla_{\text{ad}}$
17. Which of these energy transport mechanisms is *not* always operational in any given part of a star?
- (a) Conduction
  - (b) Radiation
  - (c) Convection [\*]
18. What happens when a star exceeds the Eddington limit at its surface?
- (a) The star collapses to a neutron star.
  - (b) The star undergoes a supernova.
  - (c) Convection sets in.
  - (d) A magnetic field is generated.
  - (e) Radiation pressure blows the surface layers off in a wind. [\*]
19. Which of these expressions is a criterion for convection to begin?
- (a)  $\nabla_{\text{ad}} > \nabla$
  - (b)  $\nabla_{\text{rad}} > \nabla_{\text{ad}}$  [\*]
  - (c)  $\Gamma < 1$
  - (d)  $\frac{d \ln T}{d \ln P} > 1$
  - (e)  $\frac{T}{\rho^{2/3}} < 1200$

20. Which of the following elements has the largest binding energy per nucleon?
- (a)  ${}_{26}^{56}\text{Fe}$  [\*]
  - (b)  ${}_2^4\text{He}$
  - (c)  ${}_{92}^{238}\text{U}$
  - (d)  ${}_6^{12}\text{C}$
  - (e)  ${}_{28}^{56}\text{Ni}$
21. What do the CNO cycle and the PP chain have in common?
- (a) Fraction of energy released as neutrinos
  - (b) Rate of production of  ${}_2^4\text{He}$
  - (c) Total rest mass / energy produced per  ${}_2^4\text{He}$  created [\*]
  - (d) Sensitivity to temperature
  - (e) Dependence on metallicity
22. According to the Vogt-Russell theorem, which two parameters uniquely determine the structure and evolution of a star?
- (a) Radius and effective temperature
  - (b) Composition and radius
  - (c) Luminosity and effective temperature
  - (d) Mass and composition [\*]
  - (e) Mass and radius
23. Put the components of the solar atmosphere in the correct in-out order
- (a) Chromosphere–corona–photosphere
  - (b) Photosphere–chromosphere–corona [\*]
  - (c) Corona–photosphere–chromosphere
  - (d) Photosphere–corona–chromosphere
  - (e) Chromosphere–photosphere–corona
24. What do the cores of low-mass and high-mass main sequence stars have in common?
- (a) Both have a uniform composition
  - (b) Both are burning hydrogen [\*]
  - (c) Both are convective
  - (d) Both are radiative
  - (e) Both are contracting
25. Which of these nuclear reactions doesn't occur during the pre-white dwarf evolution of a  $1 M_{\odot}$  star?
- (a) PP chain
  - (b) CNO cycle
  - (c) Carbon burning [\*]
  - (d) Triple alpha
26. What element is enriched by incomplete CNO-cycle burning?
- (a) Hydrogen
  - (b) Neon
  - (c) Carbon
  - (d) Nitrogen [\*]

- (e) Oxygen
27. What is the reason why a  $10 M_{\odot}$  main-sequence star has a convective core?
- (a) The opacity is large in the core
  - (b) The opacity is small in the core
  - (c) The nuclear energy generation has a high temperature sensitivity [\*]
  - (d) There are composition gradients in the core
  - (e) The core is contracting
28. Which of these is the principal element produced during a helium flash?
- (a) Hydrogen
  - (b) Helium
  - (c) Carbon [\*]
  - (d) Oxygen
  - (e) Silicon
29. In what evolutionary stage of a star would you expect to find an isothermal core surrounded by a hydrogen-burning shell?
- (a) After the helium flash
  - (b) Shortly after the end of the main sequence [\*]
  - (c) At the beginning of the main sequence
  - (d) On the asymptotic giant branch
  - (e) Toward the end of the main sequence
30. Where in a  $1 M_{\odot}$  star would you expect to find the CNO cycle occurring?
- (a) Core hydrogen burning
  - (b) Core helium burning
  - (c) Core carbon burning
  - (d) Shell hydrogen burning [\*]
  - (e) Shell helium burning
31. Stars above  $\approx 1.1 M_{\odot}$  evolve briefly toward higher effective temperatures at the end of their main-sequence lifetimes. What process is responsible for this blueward evolution?
- (a) Hydrogen shell ignition
  - (b) Helium shell ignition
  - (c) Overall Kelvin-Helmholtz contraction [\*]
  - (d) Onset of convection
  - (e) Core degeneracy
32. What does the Schönberg-Chandrasekhar limit correspond to?
- (a) The maximum central temperature that a convective core can have
  - (b) The maximum mean density that a non-degenerate core can have
  - (c) The maximum boundary pressure that an isothermal core can have [\*]
  - (d) The maximum total mass that a white dwarf can have
  - (e) The maximum core mass that a neutron star can have
33. What opacity source is responsible for the Hayashi line?
- (a) Bound-free

- (b) Bound-bound
  - (c) Electron scattering
  - (d) Free-free
  - (e)  $H^-$  [\*]
34. What is always true of a star lying on the Hayashi line?
- (a) It is fully convective [\*]
  - (b) it is fully radiative
  - (c) It has core helium burning occurring
  - (d) It has shell hydrogen burning occurring
  - (e) It is losing mass
35. What physical conditions are most favorable to electron degeneracy?
- (a) Low temperature and high density [\*]
  - (b) High temperature and low density
  - (c) High temperature and high density
  - (d) Low temperature and low density
36. What quantum-mechanical principle is responsible for electron degeneracy?
- (a) The Heisenberg uncertainty principle
  - (b) Schrodinger's cat
  - (c) The Pauli exclusion principle [\*]
  - (d) Wigner's friend
  - (e) The Einstein-Podolski-Rosen paradox
37. What is the cause of first dredge up?
- (a) Convective mixing on the asymptotic giant branch
  - (b) Convective mixing on the red giant branch [\*]
  - (c) Radiative levitation on the horizontal branch
  - (d) Mass loss on the main sequence
  - (e) Thermal pulses on the asymptotic giant branch
38. What element is typically brought to the surface during third dredge up?
- (a) Silicon
  - (b) Carbon [\*]
  - (c) Nitrogen
  - (d) Oxygen
  - (e) Neon
39. Between which two evolutionary stages (in order) does the helium flash occur?
- (a) Main sequence, red giant branch
  - (b) Asymptotic giant branch, red giant branch
  - (c) Horizontal branch, main sequence
  - (d) Red giant branch, horizontal branch [\*]
  - (e) Asymptotic giant branch, horizontal branch
40. Place the evolutionary stages in the correct order
- (a) Main sequence – red giant branch – horizontal branch – asymptotic giant branch [\*]

- (b) Horizontal branch – main sequence – asymptotic giant branch – red giant branch
  - (c) Asymptotic giant branch – main sequence – red giant branch – horizontal branch
  - (d) Main sequence – horizontal branch – red giant branch – asymptotic giant branch
  - (e) Red giant branch – main sequence – horizontal branch – asymptotic giant branch
41. In the core of a star above  $10 M_{\odot}$ , what happens after nuclear burning reaches  ${}^{56}_{26}\text{Fe}$  ?
- (a) It slowly cools off
  - (b) It collapses [\*]
  - (c) It explodes
  - (d) It becomes isothermal due to conduction
  - (e) It burns the  ${}^{56}_{26}\text{Fe}$  to make  ${}^{56}_{28}\text{Ni}$
42. What sorts of nuclei are produced during *r*-process nucleosynthesis?
- (a) Neutron-rich [\*]
  - (b) Electron-rich
  - (c) Low-*A*
  - (d) Proton-rich
  - (e) Low-*Z*
43. What element is always seen in the spectrum of a type II supernova?
- (a) Helium
  - (b) Silicon
  - (c) Hydrogen [\*]
  - (d) Carbon
  - (e) Oxygen
44. What phenomenon precedes complete envelope ejection during the final AGB phase of a low-mass star?
- (a) Core collapse
  - (b) Thermal pulses [\*]
  - (c) Carbon ignition
  - (d) Core helium flash
  - (e) Neutronization
45. During core-collapse in a high-mass star, what fraction of the energy is released as neutrinos?
- (a) 1%
  - (b) 10%
  - (c) 50%
  - (d) 90%
  - (e) 99% [\*]
46. What sort of star will the central star of a planetary nebula cool down to become?
- (a) White dwarf [\*]
  - (b) Brown dwarf
  - (c) Red dwarf
  - (d) Neutron star
  - (e) Black hole

47. Why is conduction so efficient in white dwarfs?
- (a) Lack of hydrogen/helium
  - (b) Electron degeneracy [\*]
  - (c) High density
  - (d) High temperature
  - (e) Lack of photons
48. How are type Ia supernova most likely formed?
- (a) Collapse of an accreting white dwarf
  - (b) Collapse of a massive star
  - (c) Merger of two black holes
  - (d) Degenerate C/O ignition in an accreting white dwarf [\*]
  - (e) Explosion of a quark star
49. What is the typical radius of a neutron star?
- (a) 1 km
  - (b) 10 km [\*]
  - (c) 100 km
  - (d) 1,000 km
  - (e) 10,000 km
50. Why don't free neutrons decay in neutron stars?
- (a) The temperature is too low
  - (b) The strong nuclear force inhibits the decay
  - (c) The decay is endothermic
  - (d) All of the possible electron states are already occupied [\*]
  - (e) The density is not high enough